

## HOMEWORK 1

Do at least 5 questions. Due September 30 at 11:59pm.

1. Let  $d \in \mathbb{Z}$  be square free and  $\neq 0, 1$ . Show that the integral closure of  $\mathbb{Z}$  in  $\mathbb{Q}(\sqrt{d})$  is  $\mathbb{Z}[\sqrt{d}]$  if  $d \not\equiv 1 \pmod{4}$  and is  $\mathbb{Z}[\frac{1+\sqrt{d}}{2}]$  if  $d \equiv 1 \pmod{4}$ .
2. Let  $A \subseteq B$  be integral domains with  $B$  integral over  $A$ . Prove that  $B$  is a field if and only if  $A$  is a field.
3. Let  $n \geq 2$  and let  $\zeta, \zeta'$  be primitive  $n$ th roots of unity in some field extension of  $\mathbb{Q}$ .
  - (a) Show that  $\frac{1-\zeta'}{1-\zeta}$  is an algebraic integer.
  - (b) Show that if  $n$  has at least two prime factors, then  $1 - \zeta$  is a unit in  $\mathbb{Z}[\zeta]$ .
4. Let  $A$  be a UFD with fraction field  $F$  and let  $E/F$  be an extension of fields. Let  $x \in E$  be algebraic over  $F$  with minimal polynomial  $f \in F[t]$ . Prove that  $x$  is integral over  $A$  if and only if  $f \in A[t]$ .
5. It can be shown that the ring of integers of  $\mathbb{Q}(\sqrt[3]{2})$  is  $\mathbb{Z}[\sqrt[3]{2}]$ . Compute the discriminant of  $\mathbb{Z}[\sqrt[3]{2}]$ .
6. Let  $F$  be a number field of degree  $n$  over  $\mathbb{Q}$  such that  $\mathcal{O}_F = \mathbb{Z}[\alpha]$  for some  $\alpha \in F$ . (The basis  $\{1, \alpha, \dots, \alpha^{n-1}\}$  is usually referred to as a *power basis* for  $F$ . Power bases don't always exist.) Let  $f$  be the minimal polynomial of  $\alpha$  over  $\mathbb{Q}$ , and let  $\alpha = \alpha_1, \dots, \alpha_n$  be the roots of  $f$ . Show that the discriminant of  $F$  equals the discriminant of  $f$ , i.e.

$$d_F = \prod_{i < j} (\alpha_i - \alpha_j)^2.$$

7. Let  $F(\alpha)/F$  be a finite separable extension of degree  $n$  generated by  $\alpha$ , let  $f \in F[t]$  be the minimal polynomial of  $\alpha$  over  $F$ , and let  $f'$  be its derivative. Show that

$$d(1, \alpha, \dots, \alpha^{n-1}) = (-1)^{\frac{n(n-1)}{2}} \text{Nm}_{F(\alpha)/F}(f'(\alpha)).$$

8. Let  $A$  be a normal Noetherian domain with fraction field  $F$ . Let  $E/F$  be a finite separable extension and let  $B$  be the integral closure of  $A$  in  $E$ .
  - (a) Let  $M \subset E$  be a finitely generated nonzero  $B$ -submodule of  $E$ . Prove that

$$M^* := \{x \in E : \text{Tr}_{E/F}(xM) \subseteq A\}$$

is also finitely generated  $B$ -submodule of  $E$ .

- (b) Consider the case of  $A = \mathbb{Z}$  and  $M = \mathcal{O}_E$  the ring of integers in a number field  $E$ . Show that  $\mathfrak{D}_{E/\mathbb{Q}} := \{x \in E : x\mathcal{O}_E^* \subseteq \mathcal{O}_E\}$  is an ideal in  $\mathcal{O}_E$ . This is called the *different* of the extension  $E/\mathbb{Q}$ .
9. Let  $E/\mathbb{Q}$  be a quadratic extension. We use the notation and definitions of Question 8.
  - (a) Compute  $\mathcal{O}_E^*$ .
  - (b) Compute the different  $\mathfrak{D}_{E/\mathbb{Q}}$  of the extension  $E/\mathbb{Q}$ .

- (c) Compute the ideal in  $\mathbb{Z}$  generated by  $\{\text{Nm}_{E/\mathbb{Q}}(x) : x \in \mathcal{D}_{E/\mathbb{Q}}\}$ . Where have you seen this before?
10. Let  $d \neq 0, 1$  be a squarefree integer, let  $F = \mathbb{Q}(\sqrt{d})$ , and let  $p$  be a prime number such that  $p \nmid 2d$ . Prove that  $p\mathcal{O}_F$  is a prime ideal in  $\mathcal{O}_F$  if and only if  $x^2 \equiv d \pmod{p}$  has no solutions in  $x \in \mathbb{Z}$ . (Hint: Note that  $\mathcal{O}_F/p\mathcal{O}_F$  is 2-dimensional over  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$ .)
11. Prove that a Dedekind domain with only finitely many prime ideals is a principal ideal domain.
12. Let  $A$  be a Dedekind domain.
- Let  $J \subseteq I$  be nonzero ideals in  $A$ . Prove there is  $a \in I$  such that  $I = J + (a)$ .
  - Prove that any ideal in  $A$  can be generated by at most two elements.
13. Prove that a Dedekind domain is a UFD if and only if it is a PID.
14. Let  $A$  be a Dedekind domain.
- Prove that for any ideals  $J \subseteq I$  of  $A$ , there is an ideal  $H$  of  $A$  such that  $J = IH$ .
  - Prove that for any nonzero ideal  $I$  of  $A$ , there is a nonzero ideal  $H$  of  $A$  such that  $IH$  is principal.
15. Let  $I$  be an ideal of a Dedekind domain  $A$ . Prove that  $I$  is a direct summand of  $A^2$  as an  $A$ -module. (Hint: Question 12. above shows there is a surjection  $f : A^2 \rightarrow I$ . To show that  $I$  is a direct summand of  $A^2$ , it suffices to show there is an  $A$ -module map  $s : I \rightarrow A^2$  such that  $f \circ s = \text{id}$ . Question 14. is useful for constructing  $s$ .)
16. Let  $A$  be a Dedekind domain and let  $S$  be a finite set of nonzero prime ideals of  $A$ . Prove that any element of  $\text{Cl}(A)$  can be represented by an ideal of  $A$  that is not divisible by any element in  $S$ .